


Patient Information Leaflet



ST MARK'S

HOSPITAL
FOUNDATION

Sphincter Exercises for people with
Bowel Control Problems


Sphincter Exercises for people with Bowel Control Problems

Introduction

Sphincter exercises can help you to improve your bowel control. When done correctly, these exercises can build up and strengthen the muscles to help you to hold both gas and stool in the back passage.

Where are these muscles?

The back passage, or anus, has two rings of muscle around it (see diagram).




The inner ring is the *internal sphincter*, an involuntary muscle, which should be closed at all times, except when you are actually trying to open your bowels. This is normally automatic, you don't have to think about doing it.

The outer ring of muscle is the *external sphincter*, which is a voluntary muscle which you can tighten up to close it more firmly if you have urgency or diarrhoea.

Both muscles wrap right around the anus. When a stool comes into the rectum the internal sphincter relaxes and allows the stool to enter the top part of the anus. Very sensitive nerves in the anal canal can tell you if it is gas or stool waiting to come out. If it is stool, you squeeze your external sphincter to stop it from coming straight out. This

squeezing moves the stool back into the rectum, where it waits until you get to the toilet.



Either or both of these sphincter muscles can become weak. This may be because of childbirth, constipation and straining or general wear and tear. Sometimes there is no obvious reason why. If you have weak muscles and cannot squeeze enough to hang on, you will feel urgency and may leak gas, liquid or even solid stool.

How can exercises help?

Exercises can strengthen these muscles so that they once again give support. This will improve your bowel control and improve or stop leakage of gas or stool. Like any other muscles in the body, the more you use and exercise them, the stronger the sphincter muscles will be.

Learning to do the exercises

It is important to learn to do the exercises in the right way, and to check from time to time that you are still doing them correctly.

Sit comfortably with your knees slightly apart. Now imagine that you are trying to stop yourself passing wind from the bowel. To do this you must squeeze the muscle around the back passage. Try squeezing and lifting that muscle as tightly as you can, as if you are really worried that you are about to leak. You should be able to feel the muscle move. Your buttocks, tummy and legs should not move at all. You should be aware of the skin around the back passage tightening and being pulled up and away from your chair. Really try to feel this. You are now exercising your anal sphincter. You should not need to hold your breath when you tighten the muscles!

Now imagine that the sphincter muscle is a lift. When you squeeze as tightly as you can your lift goes up to the fourth floor. But you cannot hold it there for very long, and it will not get you safely to the toilet as it will get tired very quickly. So now squeeze more

gently, take your lift only up to the second floor. Feel how much longer you can hold it than at the maximum squeeze.

Practising your exercises

1. Sit, stand or lie with your knees slightly apart. Slowly tighten and pull up the sphincter muscles as tightly as you can. Hold tightened for at least five seconds, then relax for about 4 seconds.

Repeat five times. This will work on the *strength* of your muscles.

2. Next, pull the muscles up to about half of their maximum squeeze. See how long you can hold this for. Then relax for at least 10 seconds.

Repeat twice. This will work on the *endurance* or staying power of your muscles.

3. Pull up the muscles as quickly and tightly as you can and then relax and then pull up again, and see how many times you can do this before you get tired. Try for about five quick pull-ups.

4. Do these exercises - five as hard as you can, two as long as you can and five quick pull-ups – four to six times every day.

5. As the muscles get stronger, you will find that you can hold for longer than five seconds, and that you can do more pull-ups each time without the muscle getting tired.

6. It takes time for exercise to make muscle stronger. You may need to exercise regularly for several months before the muscles gain their full strength.

Tips to help you

- At first it is probably a good idea to set aside some time for these exercises and really concentrate on getting them right. But quite soon they should become easy to do wherever you are. Nobody need know what you are doing!
- Get into the habit of doing your exercises with things you do regularly - every time you touch water if you are a housewife, every time you answer the phone if you are at the office whatever you do often.
- If you are unsure that you are exercising the right muscle, put a finger on the anus as you squeeze to check. You should feel a gentle lift and squeeze if you are exercising the right muscle. Or look at the area in a mirror - you should see the anus pucker up as you squeeze it.
- Use your muscles when you need them - pull up the muscles if you feel urgency and that you are about to leak. But remember that you cannot hold your tightest squeeze for very long, so you are better to use a gentler squeeze that you can hold for longer. Your control will gradually improve.

- Watch your weight - extra weight puts extra strain on your muscles.
- Once you have regained control of your bowel, don't forget your exercises. Continue to do them a few times each day to ensure that the problem does not come back.

Remember - you can do these exercises wherever you are - nobody need know what you are doing!

Do you have any questions?

This information sheet is designed to teach you how to exercise your anal sphincter muscle to improve your bowel control. If you have problems doing the exercises, or if you don't understand any part of this information sheet, ask you doctor, nurse or physiotherapist for help.

Contact details:

St Mark's Hospital, Watford Road, Harrow, Middlesex HA1 3UJ

Frederick Salmon Ward South - 020 8235 4022

Frederick Salmon Ward North – 020 8235 4191

Robert and Lisa Sainsbury Wing - 020 8869 3399

Call centre – 020 8235 4061

www.stmarkshospital.org.uk

Author: St Mark's Hospital nursing team
Date for review: January 2015